Treatment of OFF Episodes in Parkinson's Disease: An Evaluation of Physician Practices

Eric J. Pappert, MD;¹ Albert Agro, PhD¹; Nigel deGruyther, BScPharm, MBA¹ Cynapsus Therapeutics, Toronto, Ontario, Canada

CYNAPSUS

BACKGROUND

OFF episodes in PD have a significant negative impact on QoL of patients

OBJECTIVE

Determine physician practices in the treatment of OFF episodes in patients with Parkinson's disease.

METHODS

A survey was sent to 405 physicians who treat patients with PD


- 175 primary care physicians [PCP],
- 97 neurologists [NEURO] and
- 36 movement disorder specialists [MOV]).

The survey queried the following areas regarding physician practices and attitudes regarding the treatment of OFF episodes


SURVEY RESULTS


102 surveys: 50 Primary Care Physicians (PCP), 30 Neurologists (NEURO) and 22 Movement Disorder Specialists (MOV)


Respondent Demographics	All N=102	PCP N=50	NEURO and MOV N=52
# of Yrs Treating PD since Residence, mean	16.7	16.3	17.1
# of PD patients/month, mean	75.4	46.2	103.3
# of Levodopa Rx/month, mean	63.6	40.1	86.3




SURVEY RESULTS (continued)








SURVEY RESULTS (continued)





CONCLUSIONS

- Physicians are concerned about OFF episodes.
- Physicians feel that their patients are aware of OFF episodes.
- Physicians feel that there is a fairly high level of unmet need regarding treatment options for OFF episodes.

ACKNOWLEDGEMENTS/DISCLOSURES

This study was supported by Cynapsus Therapeutics. EJP, ND, and AA are all employees of Cynapsus Therapeutics and hold stock or stock options.